Information for breastfeeding families

Calming a Colicky Breastfed Baby



Crying is often related to hunger, and feeding is the obvious solution. However, periods of crying, with no obvious cause and no ready cure, plague many parents. Here are a few suggestions. Give several a try. One may work today and another may work tomorrow. Be flexible, and know that "this too will pass".

Symptoms:

Periods of inconsolable crying Clenching fists Drawing legs up to abdomen Stiffening as if in pain

Passing gas Grimacing Possible causes: (may be more than one)

Immature gastro-intestinal system

Sensitive nervous system

Excessive gas or over-feeding

Formula intolerance

Reaction to foods the breastfeeding

mother has eaten Over-stimulation

Possible remedies:

- ✓ Hold your baby skin-to-skin several times each day
- ✓ Motion in any form

Walking

Baby swing

Rocking in rocking chair

Rocking side to side while standing

Ride in the car or in a stroller

Carry baby in infant sling

- ✓ Over-the-counter gas-relief drops (discuss with MD)
- ✓ Singing, humming
- ✓ Take a break, let someone else try for awhile; babies sense your tension
- √ White sound nearby (running water, vacuum, clothes dryer, hair dryer)
- √ Commercially available recordings with strong beat designed to simulate intrauterine sounds.
- ✓ Swaddle your baby snugly when crying
- ✓ Undress your baby and allow complete freedom of motion
- ✓ Avoid over-stimulation from noises, lights or motion
- ✓ Distract the baby with different sounds, sights or places
- ✓ Place pressure on your baby's abdomen (gently)

Hold baby in the "colic hold" (facing floor, supported by your arm, heel of your hand putting pressure on the abdomen)

Hold baby over your shoulder or over your knees

- ✓ Bicycle your baby's legs
- ✓ Take an infant massage class
- ✓ Keep a food diary to determine if a particular food bothers your baby (takes 4-6 hours for most foods to reach baby)
- ✓ Consult a Lactation Consultant. She may suggest feeding on only one breast per feeding or other techniques to balance the "foremilk" and "hindmilk" your baby obtains while feeding
- ✓ Contact your physician for other medical explanations and solutions



"Colic hold"